References:

Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, Filiberti A, Flechtner H, Fleishman SB, de Haes JCJM, Kaasa S, Klee MC, Osoba D, Razavi D, Rofe PB, Schraub S, Sneeuw KCA, Sullivan M, Takeda F, The European Organisation for Research and Treatment of Cancer QLQ-C30: A quality-of-life instrument for use in international clinical trials in oncology: Journal of the National Cancer Institute, 1993; 85: 365-376.

Alexandre J, Gross-Goupil M, Falissard B, Nguyen ML, Gornet JM, Misset JL, Goldwasser F, Evaluation of the nutritional and inflammatory status in cancer patients for the risk assessment of severe haematological toxicity following chemotherapy: Ann Oncol, 2003;14(1):36-41.

Argiles JM, Busquets S, Felipe A, López-Soriano FJ, Muscle wasting in cancer and ageing: cachexia versus sarcopenia: Adv Gerontol, 2006;18:39-54.

Arrieta O, Ortega RMM, Villanueva-Rodríguez G, Sema-Thome MG, Flores-Estrada D, Diaz-Romero C, Rodríguez CM, Martínez L, Sanchez-Lara K, Association of nutritional status and serum albumin levels with development of toxicity in patients with advanced non-small cell lung cancer treated with paclitaxel-cisplatin chemotherapy: a prospective study: BMC Cancer, 2010, 10:50

Baracos VE, Reiman T, Mourtzakis M, Gioulbasanis I, Antoun S, Body composition in patients with non-small cell lung cancer: a contemporary view of cancer cachexia with the use of computed tomography image analysis: Am J Clin Nutr, 2010;91(4):1133S-1137S.

Bauer J, Capra S, Ferguson M, Use of the scored Patient-Generated Subjective Global Assessment (PG-SGA) as a nutrition assessment tool in patients with cancer: European journal of clinical nutrition, 2002; 56(8):779-85.

Bohannon RW, Dynamometer measurements of hand-grip strength predict multiple outcomes: Percept Motor Skills, 2001; 93:323–8.

Boker B, Lüders H, Grohé C, Prognostic Relevance of Body Mass Index and Rash for Patients with Metastatic Non-Small-Cell Lung Cancer under Therapy with Erlotinib": Pneumologie, 2012;66(2):89-95.

Chandrasekaran B, Ghosh A, Prasad C, Krishnan K, Chandrasharma B, Age and anthropometric traits predict handgrip strength in healthy normals: J Hand Microsurg, 2010;2(2):58-61.

Detsky AS, Baker JP, Mendelson RA, Wolman SL, Wesson DE, Jeejeebhoy KN, Evaluating the accuracy of nutritional assessment techniques applied to hospitalized patients: methodology and comparisons: Jpen, 1984; 8(2):153-9.

Dewys WD, Begg C, Lavin PT, Band PR, Bennett JM, Bertino JR, Cohen MH, Douglass HO Jr, Engstrom PF, Ezdinli EZ, Horton J, Johnson GJ, Moertel CG,Oken MM, Perlia C, Rosenbaum C, Silverstein MN, Skeel RT, Sponzo RW, Tormey DC, Prognostic effect of weight loss prior to chemotherapy in cancer patients. Eastern Cooperative Oncology Group: Am J Med, 1980;69(4):491-7.

Fried D, Lawrence M, Khandani AH, Rosenman J, Cullip T, Chera BS, Is Image Registration of Fluorodeoxyglucose-Positron Emission Tomography/ComputedTomography for Head-and-Neck Cancer Treatment Planning Necessary?: Int J Radiat Oncol Biol Phys, 2012.

Gioulbasanis I, Baracos VE, Giannousi Z, Xyrafas A, Martin L, Georgoulias V, Mavroudis D, Baseline nutritional evaluation in metastatic lung cancer patients: Mini Nutritional Assessment versus weight loss history: Ann Oncol, 2011;22(4):835-41.

Gordon JN, Green SR, Goggin PM, Cancer cachexia: Qjm, 2005; 98(11):779-88.

Harvie MN, Howell A, Thatcher N, Baildam A, Campbell,
Energy balance in patients with advanced NSCLC, metastatic melanoma and metastatic breast cancer receiving chemotherapy- a longitudinal study: Br J Cancer, 2005; 28;92(4):673-80).

Hebert JR, Augustine A, Barone J, Kabat GC, Kinne DW, Wynder EL, Weight, height and body mass index in the prognosis of breast cancer: early results of a prospective study: Int J Cancer, 1988; 15;42(3):315-8.

Hill GL, Jonathan E. Rhoads Lecture, Body composition research- implications for the practice of clinical nutrition: J Parenter Enteral Nutr, 1992;16:197–218.

Hurria A, Togawa K, Mohile SG, Owusu C, Klepin HD, Gross CP, Lichtman SM, Gajra A, Bhatia S, Katheria V, Klapper S, Hansen K, Ramani R, Lachs M, Wong FL, Tew WP, Predicting chemotherapy toxicity in older adults with cancer: a prospective multicenter study: J Clin Oncol, 2011; 1;29(25):3457-65.

Humphreys J, de la Maza P, Hirsch S, Barrera G, Gattas V,Bunout D, Muscle strength as a predictor of loss of functional status in hospitalised patients: Nutrition, 2002;18:616–20.

Hunt DR, Rowlands BJ, Johnston D, Hand grip strength— a simple prognostic indicator in surgical patients: J Parenter Enteral Nutr, 1985;9:701–14.

Hurria A, Hurria A, Zuckerman E, Panageas KS, Fornier M, D'Andrea G, Dang C, Moasser M, Robson M, Seidman A, Currie V, VanPoznak C, Theodoulou M,Lachs MS, Hudis C, A prospective, longitudinal study of the functional status and quality of life of older patients with breast cancer receiving adjuvant chemotherapy: J Am Geriatr Soc, 2006;54(7):1119-24.

Iversen PO, Wisløff F, Gulbrandsen N, Reduced nutritional status among multiple myeloma patients during treatment with high-dose chemotherapy and autologous stem cell support: Clinical Nutrition, 2010;29(4):488-91. Epub 2009; 30.

Jeejeebhoy KN. Bulk or bounce—the object of nutritional support. J Parenter Enteral Nutr 1988;12:539–49.

Keys A, Brozek J, Henschel A, Mickelsen O, Taylor HF, The biology of human starvation. Minneapolis: University of Minnesota Press, 1950.

Klidjian AM, Foster KJ, Kammerling RM, Cooper A, Karran SJ, Relation of anthropometric and dynamometric variables to serious postoperative complications: BMJ, 1980;281: 899–901.

Klopp AH, Eifel PJ, Biological predictors of cervical cancer response to radiation therapy: Semin Radiat Oncolology, 2012 ;22(2):143-50.

Kuzuya M, Kanda S, Koike T, Suzuki Y, Satake S, Iguchi A, Evaluation of Mini-Nutritional Assessment for Japanese frail elderly: Nutrition (Burbank, Los Angeles County, Calif), 2005; 21(4):498-503.

Laky B, Janda M, Bauer J, Vavra C, Cleghorn G, Obermair A, Malnutrition among gynaecological cancer patients: European journal of clinical nutrition, 2007; 61(5):642-6.

Leitch EF, Chakrabarti M, Crozier J E M, McKee R F, Anderson J H, Horgan P G, McMillan D C, Comparison of the prognostic value of selected markers of the systemic inflammatory response in patients with colorectal cancer: Br J Cancer, 2007; 5; 97(9): 1266–1270.

Lobo DN, Allison SP, Nutritional support and functional recovery: Curr Opin Clin Nutr Metab Care, 2000;3:129–34.

Lochs H, Dervenis C, Malnutrition–the ignored risk factor: Digestive diseases (Basel, Switzerland), 2003; 21(3):196-7.

McMillan DC, Systemic inflammation, nutritional status and survival in patients with cancer: Curr Opin Clin Nutr Metab Care, 2009; 12(3):223-6.

Mohan A, Singh P, Kumar S, Mohan C, Pathak AK, Pandey RM, Guleria R, Effect of change in symptoms, respiratory status, nutritional profile and quality of life on response to treatment for advanced non-small cell lung cancer: Asian Pac J Cancer Prev, 2008; 9(4):557-62.

Mrian M, Robert S, Clinical Nutrition for Oncology patients: Jones and Bartlett, 2010,1.

Muscaritoli M, Bossola M, Aversa Z, Bellantone R, Rossi Fanelli F, Prevention and treatment of cancer cachexia- new insights into an old problem: Eur J Cancer, 2006; 42(1):31-41.

Oh CA, Kim DH, Oh SJ, Choi MG, Noh JH, Sohn TS, Bae JM, Kim S, Nutritional risk index as a predictor of postoperative wound complications after gastrectomy: World J Gastroenterol, 2012;18(7):673-8.

Ottery FD, Cancer cachexia- prevention, early diagnosis, and management: Cancer practice, 1994, 2(2):123-31.

Perdue C, Understanding nausea and vomiting in advanced cancer: Nurs Times, 2005; 25-31;101(4):32-5.

Phippen NT, Lowery WJ, Barnett JC, Hall LA, Landt C, Leath CA, Evaluation of the Patient-Generated Subjective Global Assessment (PG-SGA) as a predictor of febrile neutropenia in gynecologic cancer patients receiving combination chemotherapy- a pilot study: Gynecol Oncol, 2011;123(2):360-4.

Prado CM, Baracos VE, McCargar LJ, Reiman T, Mourtzakis M, Tonkin K, Mackey JR, Koski S, Pituskin E, Sawyer MB, Sarcopenia as a determinant of chemotherapy toxicity and time to tumor progression in metastatic breast cancer patients receiving capecitabine treatment: Clin Cancer Res., 2009; 15;15(8):2920-6.

Romero-Ventosa EY, Mucientes-Molina A, Pedrido-Reino E, Lago-Rivero N, Constenla-Caramés L, Arias-Santos I, Efficacy and toxicity of erlotinib in non-small cell lung cancer treatment: Farm Hosp, 2012;36(2):68-76.

Ryu SW, Kim IH, Comparison of different nutritional assessments in detecting malnutrition among gastric cancer patients: World J Gastroenterol, 2010 ;14;16(26):3310-7.

Sarraf KM, Belcher E, Raevsky E, Nicholson AG, Goldstraw P, Lim E, Neutrophil/lymphocyte ratio and its association with survival after complete resection in non-small cell lung cancer: J Thorac Cardiovasc Surg, 2009; 137(2):425-8.

Sawada NO, de Paula JM, Sonobe HM, Zago MM, Guerrero GP, Nicolussi AC, Depression, fatigue, and health-related quality of life in head and neck cancer patients- a prospective pilot study: Support Care Cancer, 2012

Smets EM, Garssen B, Schuster- Uuitterhoeve AL, de Haes JC, Fatigue in cancer patients: Br J Cancer, 1993; 68, 220-224.

Smith RA, Bosonnet L, Raraty M, Sutton R, Neoptolemos JP, Ghaneh P, Preoperative platelet-lymphocyte ratio is an independent significant prognostic marker in resected pancreatic ductal adenocarcinoma: Am J Surg, 2009; 197(4):466-72.

Stratton RJ, Green CJ, Elia M, Disease related malnutrition— an evidence-based approach to treatment: Oxford CAB International, 2003.

Urba S, Gatz J, Shen W, Hossain A, Winfree K, Koustenis A, Peterson P, Cohen EE, Quality of life scores as prognostic factors of overall survival in advanced head and neck cancer- Analysis of a phase III randomized trial of pemetrexed plus cisplatin versus cisplatin monotherapy: Oral Oncolology, 2012

Vassallo JABE, Actualización Ponderada de los Factores de Riesgo del Cancer. Montevideo: Comisión Honoraria de Lucha contra el Cáncer 2003.

Walsh SR, Cook EJ, Goulder F, Justin TA, Keeling NJ, Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer: J Surg Oncol, 2005, 91(3):181-4.

Webb AR, Newman LA, Taylor M, Keogh JB, Hand grip dynamometry as a predictor of postoperative complications reappraisal using age standardized grip strengths: J Parenter Enteral Nutr, 1989; 13:30–3.

Yamanaka T, Matsumoto S, Teramukai S, Ishiwata R, Nagai Y, Fukushima M, The baseline ratio of neutrophils to lymphocytes is associated with patient prognosis in advanced gastric cancer: Oncology, 2007; 73(3- 4):215-20, Epub 2008; 17.